



Recombinant Human Gamma-adducin (ADD3)

Product Code	CSB-MP887015HU
Abbreviation	ADD3
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q9UEY8
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	>85% (SDS-PAGE)
Sequence	SSDASQGV I TTPPPSPMPH KERYFDRINE NDPEYIRERN MSPDLRQDFN MMEQRKRVTQ ILQSPAFRED LECLIQEQMK KGHNP TGLLA LQQIADYIMA NSFSGFSSPP LSLGMVTPIN DLPGADTSSY VKGEKLTRCK LASLYRLVDL FGWAHLANTY ISVRISKEQD HIIIPRGLS FSEATASNLV KVNIIGEVVD QGSTNLKIDH TGFSPHAAIY STRPDVKCVI HIHTLATAAV SSMKCGILPI SQESLLLGDV AYYDYQGSLE EQEERIQLQK VLGPSCKVLV LRNHGVVALG ETLEEFHYI FNVQLACEIQ VQALAGAGGV DNLHVLDLDFQK YKAFTYTVAA SGGGGVNMGS HQKWVGEIE FEGLMRTLND LGYRTGYAYR HPLIREKPRH KSDVEIPATV TAFSFEDDTV PLSPLKYMAQ RQREKTRWL NSPNTYMKVN VPEESRNGET SPRTKITWMK AEDSSKVSGG TPIKIEDPNQ FVPLNTPNE VLEKRNKIRE QNRYDLKTAG PQSQLLAGIV VDKPPSTMQF EDDDHGPPAP PNPFSHLTEG ELEEYKRTIE RKQQGLEDAE QELLSDDASS VSQIQSQTQS PQNVPEKLEE NHELFSKFSI SMEVPMVVN GKDDMHDVED ELAKRVSRLS TSTTIENIEI TIKSPEKIEE VLSPEGSPSK SPSKKKKKFR TPSFLKKNKK KEKVEA
Source	Mammalian cell
Target Names	ADD3
Protein Names	Recommended name: Gamma-adducin Alternative name(s): Adducin-like protein 70
Expression Region	2-706
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full Length of Mature Protein
Target Details	Adducins are heteromeric proteins composed of different subunits referred to as adducin alpha, beta and gamma. The three subunits are encoded by distinct genes and belong to a family of membrane skeletal proteins involved in the assembly of spectrin-actin network in erythrocytes and at sites of cell-cell contact in epithelial tissues. While adducins alpha and gamma are ubiquitously



expressed, the expression of adducin beta is restricted to brain and hematopoietic tissues. Adducin, originally purified from human erythrocytes, was found to be a heterodimer of adducins alpha and beta. Polymorphisms resulting in amino acid substitutions in these two subunits have been associated with the regulation of blood pressure in an animal model of hypertension. Heterodimers consisting of alpha and gamma subunits have also been described. Structurally, each subunit is comprised of two distinct domains. The amino-terminal region is protease resistant and globular in shape, while the carboxy-terminal region is protease sensitive. The latter contains multiple phosphorylation sites for protein kinase C, the binding site for calmodulin, and is required for association with spectrin and actin. Alternatively spliced adducin gamma transcripts encoding different isoforms have been described. The functions of the different isoforms are not known.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.