



# Recombinant Bovine Interleukin-4 (IL4)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-MP011659BO  |
| <b>Storage</b>           | Store at -20°C, for extended storage, conserve at -20°C or -80°C.   |
| <b>Uniprot No.</b>       | P30367  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Bos taurus (Bovine)   |
| <b>Purity</b>            | >85% (SDS-PAGE)   |
| <b>Sequence</b>          | HKCDIT LAEIIKTLNI LTTRKNSCME LPVADVFAAP KNTTEKETFC<br>RVGIELRRIY RSHTCLNKFL GGLDRNLNSL ASKTCSVNEA KTSTSTLKDL<br>LERLKTIMKE KYSKC  |
| <b>Source</b>            | Mammalian cell  |
| <b>Target Names</b>      | IL4   |
| <b>Protein Names</b>     | Recommended name: Interleukin-4 Short name= IL-4 Alternative name(s): B-cell stimulatory factor 1 Short name= BSF-1 Lymphocyte stimulatory factor 1   |
| <b>Expression Region</b> | 25-135  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | Tag type will be determined during the manufacturing process.   |
| <b>Protein Length</b>    | Full Length of Mature Protein   |
| <b>Target Details</b>    | This protein is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. |
| <b>Reconstitution</b>    | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.   |
| <b>Shelf Life</b>        | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |